
## The SrTiO3 displacive transition revisited by Coherent X-ray Diffraction

## David Le Bolloc'h<sup>1</sup>, Sylvain Ravy<sup>2</sup>, Rolland Currat<sup>3</sup>, Andrei Fluerasu<sup>4</sup>, C. Mocuta<sup>4</sup>, B. Dkhil<sup>5</sup>

<sup>1</sup>Laboratoire de Physique des Solides, Bât. 510, Université Paris-Sud, 91405 Orsay, France <sup>2</sup>Synchrotron-Soleil, L'Orme des merisiers, Saint-Aubin, BP 48, 91192 Gif-sur-Yvette, France <sup>3</sup>Institut Laue Langevin, 6, rue Jules Horowitz BP 156 - 38042 Grenoble Cedex 9, France <sup>4</sup>ESRF, Grenoble, France; <sup>5</sup> LSPMS, École centrale Paris, 92295 Châtenay-Malabry, France

Although most issues concerning the application of the scaling theory of phase transitions have been settled long ago, two frequently observed scattering features remain unaccounted for within standard scaling theory: the "neutron" central peak (CP) and the "x-ray" narrow component (NC). Remarkably, both features were first evidenced in studies of the critical behavior associated with the T<sub>c</sub> ~100-105 K antiferrodistortive transition in the perovskite SrTiO<sub>3</sub>. We present a Coherent X-ray Diffraction study of the antiferrodistortive displacive transition of SrTiO<sub>3</sub>. From the microbeam x-ray coherent diffraction patterns, we show that the broad (short-length scale) and the narrow (long-length scale) components can be spatially disentangled. Moreover, both components exhibit a static speckle pattern. This gives evidence that the narrow component corresponds to static ordered domains. We interpret the speckles in the broad component as due to a very slow dynamical process, corresponding to the well-known central peak seen in inelastic neutron scattering. Beyond this experiment, we show that CXD is a valuable new tool to study phase transitions and defects in the low temperature ordering.

First results presented at Coherence 2005 and IUCr2005. To be published in Phys. Rev. Lett. (Cond-mat/0611308)



2D pattern of the (3/2,1/2,1/2) superstructure reflection at two different positions of the beam. Broad and narrow components can be disentangled by  $\mu$ -beam. On both components static speckles are observed on a ~10 mn time scale.